terested parts directly or indirectly; but I do not approve of invidious attacks made from behind the bush.—*Grower Fay, Lee, Kent.*

Early Peas.—In reference to Mr. M'Ewen's report on Peas I think he must have been wrong sort in No. 1. I have grown it every year with the Emperor since the first season in which it came out, and I have always found it about eight days earlier; in fact the whole of last June by the time the other comes in. I frequently sow from 15 to 20 acres of Peas on the same day on purpose to test them, and I find the Emperor to be a general surpass. I think Fairbeard's Conqueror will stand high as a first early. Dixon's Favourite, which is about as early as the Prince, appears to be a good Pea, producing as many as 12 Peas in a pod, which must ensure a good supply. *Richard Ritott, Donnington, near Spalding.*

The subject of Deep Wells has been sometimes discussed in your columns. I have a well 325 feet deep, and the 12-gallon bucket actually weighs 232 lbs.; this, with many yards I used a child's weighing 232 lbs.; this, with the water, itself 96 lbs., amounts to 481 lbs. I have made an enormous swing of labour by using one of the 1st half-year's Journal's rules over me. Now, will any one have the charity to say from experience whether there could not be a great saving in the weight of the bucket. Would zinc, or gutta percha, or leather serve? The bucket must be strong enough to withstand being occasionally dashed against the side of the well. I must I stick to my old substantial oaken friend?—*C. D.*

Proliferous Rose.—I send you a curious "freak of nature" in a double white Stock. At first sight it presents nothing remarkable, but having watched its development I do not remember to have seen such a curiosity before. Several others are to be found in the ordinary way; when the outer petals began to die off the centre of the flowers commenced swelling, and gradually formed little clusters of buds, as in the drawing sent. There appeared all round the under side of the lip of the saucer shaped trey from 20 to 30 nearly invisible filaments, 1 inch in length, with small white egglike specks at the extremities. They project like little bristles, vibrate with the least breath, yet adhere very firmly. The Rose is a Bourbon, Souvenir de Malmaison, or japanned, is in constant use, and dusted every morning. *P. W. M., Shirley, Croydon.* [These are evidently the stalked eggs of a Hemerobius.]

Societies.

CALEDONIAN HORTICULTURAL: *June* 27.—The following is a report of the prizes awarded on this occasion. Shrubby Greenhouse Plants: 1st, Mr. Lockhart also for R. Dundas, Esq., with Tetratheca verticillata and Boronia heterophylla. Mr. Lockhart also received the prize for Cape Heaths, the finest produced being Cavendishii and Cavendish. Two Pelargoniums: 1st, Mr. Reid, gr. to W. Wilson, Esq., with Zara and Star (Beck's); Mr. Reid, 2d, with Zara and Mrs. Dalbousie, with Attraction and Forget-me-not. Four Pelargoniums in 6-inch pots: 1st, Mr. Forrest, gr. to Messrs. Moss, with Carlo, Lady Alice, and two Wonderful; 2d, Mr. Henderson, gr. to C. K. Sivewright, Esq., with Fair Helen, Petruchio, Carlos, and Seraskier. Two Fancy Pelargoniums: 1st, Mr. Stewart, with decora and Queen Victoria: 2d, Mr. Reid, with Carlotta Grisi and Annette; 3d, Mr. Forrest, with decora and Lady Downes. Four Fancy Pelargoniums in 6-inch pots: 1st, Mr. Henderson, with crubescens, formosissimum, Celestial, and Critair; 2d, Mr. Forrest, with Madame Soutag, Madame Vaucher, Mrs. Campbell, and Annette. Two Dark Fuchsias in 8-inch pots: 1st, Mr. Walker, gr. to J. Mood, Esq., with Prince Albert and Ounega; Mr. Henderson, with Countess of Sefton and Prince Albert. Two Light Fuchsias in 8-inch pots: 1st, Mr. Walker, with Queen of Hanover and Mrs. Lee; 2d, Mr. Henderson, with Mrs. Kent and Venus de Medici. In Roses, the 1st prize was gained by M. Buist, Esq., Tynninghame, whose stand contained the following, viz., Général Jacqueminot, Baronne Prevost, Jules Margottin, Louise Peyronny, Charles Lawson, Devoniensis, Abricoté, Paul Ricaut, Reine des Fleurs, Triomphe de Rennes, Paul and General Jacqueminot. A second premium was voted to Mr. Stewart, for Great Western, Mrs. Rivers, Lanel, Elize Sauvage, Blanchfleur, Celine, Madame Knithi, Gloire de Rosamene, Baronne de Montpensier, Devoniensis, General Kleber, and Madame Humboldt. Phloxes: 1st, Mr. M'Farlane, with Barriton, Mrs. Young, Macgregor, Rival, and a seedling named Mrs. M'Call, in the way of Abdel Medschid Khan, from which it had been raised. Calceolarias: 1st, Mr. Stewart, with Falstaff, Sir Walter Scott, George Shelley, and a seedling. Several extra prizes were offered by the office-bearers, the first of which (two sovereigns) for the best collection of plants and Dicentra Pluma, was awarded to Mr. Blair, gr. to G. C. Arbuthnot, Esq. His collection included Tetratheca verticillata, Leschenaultia formosa, Epacris

minista, Mahernia incisa, Erica breviflora, centricosa superba, and tricolor flamma, Statice Hubfordi, Pimelea decussata, Hoya bella, Ixora coccinea, Oncidium flexuosum, Aphelexia macrantha purpurea, Æschynanthus speciosus, and Clerodendron Kæmpferi. Extra prizes were also placed on the tables for exhibition only were as follows—From the nursery of Messrs. Dickson & Co. were Pelargoniums, Gloriosa, &c.; from Messrs. P. Lawson & Son, Calceolarias and Heaths; from Messrs. J. Dickson & Sons, Pelargoniums, including French seedlings, &c.; from Mr. Mood, Pine, rare Alpine plants and Ferns, including Cheilanthes alabamensis, Gymnogramma peruviana, Nothochlæna nivea, Todea pellinelda, and Adiantum Moorei; from Mr. Douglas, several trays of Pelargoniums, Rose, Verbenas, &c.; from Messrs. Wright, Renwick, & Co., Dracæna rubella, Scarlet Pelargoniums, &c.; from Messrs. Downie & Laird, spikes of Delphinium formosum, Calceolarias, Petunias, Phlox Addisoni, &c.; from Mr. C. Lumsden, Pelargoniums; from Mr. Henderson, Linum grandiflorum rubrum, and new Shrubby Calceolarias; from Messrs. Cunningham, Fraser, & Co., boxes of Pelargoniums, Calceolarias, Phlox, and Roses; from Mr. Lockhart, Cissus discolor; from Mr. Stewart, Gloxinias; from Mr. Blair, Stanhopea oculata; from Mr. Buist, Bonaria Senecioides; from Mr. Forrest, and fancy Pelargoniums; from Mr. Young, Stocks; from Mr. Duncan Kerr, a seedling Pansy; from Mr. Ferguson, Dammendonea Donii; and from Mr. Dunn, Stocks. Miss Barneton, 26, Clarence Street, exhibited a neat group of wax fruit.

EDINBURGH BOTANICAL: *June* 11.—The President in the chair. The following papers were read, viz.:—1. "On the Fungi of Potatoes, Diseased and other vegetable Parasites, with Aspergillus glaucus," by Mr. John Lowe. The object of this communication was to show the relation of the parasitic fungi to the parasitic growth in Foreign fuscus and other skin diseases, and a common species of fungus, Aspergillus glaucus; and to establish the fact that a number of these forms which have hitherto been regarded as specifically distinct. After adverting to the difficulties experienced in studying the minute forms of vegetation, the author proceeded to give a detail of the experiments which had been made with the view of producing the plant in a state of perfect fructification. He then enabled to of ascertain its specific value. A quantity of flavus crust having been procured from a case of Perigo lupinosa, a portion was immersed in distilled water, another was placed on cheese, and a third in solution of raw sugar. The first of these did not germinate, but became disintegrated in a few days. This was probably owing to the temperature of the fluid not being sufficiently high, as it is well-known that the yeast plant germinates in a few hours in solutions at an elevated temperature, during the manufacture of Butyric acid. The cells placed on cheese also failed to germinate, but those in the sugar solution soon began to put into glycerine. Those immersed in the saccharine solution gave a different result. At the end of 48 hours the cells had germinated, and were found to contain in frait, on the day following they began to mute into moniliform chains forming a mycelium, the filaments of which after a time were converted into spherical sporules. At the end of about a month the perfect fructification of Aspergillus glaucus appeared. During the growth of the plant, the stages of development were carefully observed daily under the microscope, and the whole of the following species (so-called) were found accurately represented as to their structure in one or other of the forms produced. Microsporon furfur, Robin; Odium albicans, Ch. R.; Torula guttata; Trichophyton, tonsurans, Malmsten; Achorion Schönleini, Remak; Porrigo Audouini, Ch. R.; M. Mentagrophytes, Ch. R.; Achorion Schönleini Remak; Leptomitus, six species; Mucor, several species; and Botrytis Robinii. With regard to the majority of these, the author remarked that they could not be with certainty considered as distinct species, and that there was every probability of such being the case; but, from the exact identity of form; and 2d, from the circumstance that they are found only in distinct species, as shown by their never or rarely producing fruit; proving them to be mere variations of some other species produced by peculiar circumstances, and not arriving at a perfect development. With Achorion Schönleini the case is different. The following facts may be stated in support of its alleged identity with Aspergillus:—1st. The sporules of the former, carefully watched during their growth, developed the peculiar characters of the latter, while, 2dly, it produced in a state of fructification in the sir sacs of birds, showing the possibility of its growing on animal tissues. 3dly. The figures given by Dr. Bennett of a section of the scalp affected by Favus, exhibits the true fructification of an Aspergillus. The author concluded with some lengthened observations on the circumstances under which this fungus is developed, and its distribution and properties. 2. "On the Properties of Lolium temulentum." By Mr. Lowe. After noticing the physiological effects which have been ascribed to the action of Darnel, the author remarked that there exists a great want of information as to the amount of the seed sometimes to produce these results. From all that has been written on the subject, it would appear as if the virulence of the herb varied in different localities. A series of experiments was given in detail showing that Darnel grown in the Botanic Garden produced no effect when taken in doses of half an ounce daily. The experiments of Dr. Christison on the Cinanthe crocata show an analogous result, this plant being a virulent poison when grown in England, Germany, and France, while in the same latitude it is seen in the Cannabis indica, which only yields its gummi resin when grown in a hot climate. Further experiments are required with regard to Darnel. The difficulty experienced in obtaining the pure seed has prevented the above series from being extended further, but it is hoped shortly to be able to make them with seed from different parts of Scotland. 3. "Further Observations on Dust Showers." By Mr. G. Lawson. After some preliminary observations, referring to the subject treated on in the previous paper on this subject, Mr. Lawson laid before the Society a letter from Dr. J. O. Whittaw, R.N., in which that gentleman remarked:—"In your paper on dust showers you allude to the sand showers described by Humboldt and by Ehrenberg as occurring near the Cape de Verd Islands, when Captain Tuckey, navigating the ocean became covered with sand. While I was at Boa Vista, the easternmost of the Cape de Verd group, during the months of May, June, July, and part of August 1846, I had ample opportunity of witnessing these phenomena. In my meteorological register, which includes observations on the direction of the winds, state of the atmosphere, thermometer, the dry and wetted bulb thermometer, the temperature of the sea, the force and direction of the current, and the nature of the clouds, I find that in April (1846) the atmosphere is recorded as hazy, and filled with sand, ten days; in May, eleven days; in June, five days; and during the first ten days of July, three days. As a general rule, when these sand fogs prevailed, the north-east trade winds were blowing with their usual force; they sometimes continued for three or four days without any intermission. At the period of their prevalence, the sand heaps which abound in the island, from disintegration, were seen to travel about from the windward to the leeward side of the island, filling the hollows in the plains, and, sometimes in the course of a night, completely altering the line of pathways, and thus bewildering the newly arrived traveller. I was in the leeward side of the island when the first of these sand fogs made its appearance, and I differed in opinion as to its source, some saying that it came from the beach and sand hills, on the windward side of the island, while others asserted that it came from a considerable distance, and that it was the African Desert. I had soon an opportunity of ascertaining that they did not originate from the beach, for I witnessed on a sand shower of considerable density, over the sea to windward of the island, between which and the African coast I was sailing, and which I considered as came to the conclusion that that coast was its source." 4. "Analogy between the serial arrangements of the Leaves of Plants, and crystalline forms." By Dr. W. Mitchell. Mr. Mitchell remarks, "Having some time ago had my attention drawn to the series expressing the general arrangement of the leaves of plants, and more recently to the former series as regulating the shapes of cones, I was led to inquire whether a similar relation might not be found to exist in crystalline forms. The result of my observations is, that the same law prevails in crystals as in plants." Mr. Mitchell examined the forms of the cubic, pyramidal, prismatic, rhombohedral, and prismatic systems, and found that, as in plants $\frac{a}{b}, \frac{a}{c}, \frac{c}{c}$ represent any three consecutive terms of the series, such as $\frac{1}{2}, \frac{1}{3}, \frac{2}{5}$, then $\frac{a+c}{b+f}, \frac{c}{f}$ will express their relation. He concludes "may we not infer, from the serial law here shown to be common to both, that a series of nodes in plants corresponds to a series of similar crystalline forms; and thatsimilar principles of law govern both classes of bodies? At least the subject seems worthy of consideration, and might lead to still closer analogies, illustrative of unity in their diversity, which we already know to be a distinguishing characteristic of the works of God."

Notices of Books.

Memoirs of the late Rt. Hon. Sir Robert Peel. 2 vols., 8vo. Murray. Edited by Lord Mahon (now Earl Stanhope) and the Rt. Hon. E. Cardwell.

The excessively political character of these interesting memoirs precludes our even referring to their very important contents, with the exception of what concerns the Potato disease, and the repeal of the corn laws, which was the subject of the 3rd book, in the second volume. Sir Robt. Peel himself describes the great calamity that visited nearly all Europe in 1846, commencing with a letter received by Sir James Graham from the Isle of Wight on the 11th August, 1846, and ending with a letter to Lord Aberdeen dated July 15, 1847. The whole of the letters are of considerable facile bound, as one concerned in the non-political events of the time, to bear his testimony to two points: one is the entire absence of all exaggeration or conceitedness

© The Complete Work of Charles Darwin Online